

Issues of Scale and Inference

Colleen M. Handel

USGS Alaska Science Center




Boreal Partners in Flight

At what geographic scale should we monitor long-term trends?


- Ecoregion
- Land-management unit

Sample size requirements


- Number of samples needed is largely independent of the size of area being monitored




Population trend is slope of line




. . . or curve



Relative power to detect regional trend




Low




High

Triggering more intense efforts



Regional decline




Local decline



Flexibility in sampling design

- Geographically stratified random design
 - Can intensify effort within strata
 - Can sample specific land units more intensively if desired
 - Can conduct initial inventory when resources available
 - Can repeat inventory sample at some later time



Repeat of initial inventory to test local trend





Yukon-Charley



Yukon-Charley inventory




Yukon-Charley




Sample design

Specific management
question with current
landscape



Pre-stratified random
sample

Long-term monitoring for
unanticipated factors or
changing landscape



Random sample with
post-stratification

Landscape-level changes across time

- Management questions
 - Fire
 - Disease and insect outbreaks
 - Timber management
 - Resource development
 - Recreational use
 - Global warming
 - Changes in biodiversity

Flexibility in sampling design

- Can continue to refine picture of current status, distribution, and habitat use
 - Survey new additional random plots each year when resources available
 - Don't need to be repeated
 - Can be repeated at some point in future
- Can ask specific management questions with same design, though less efficient